



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

21 Robin	27 Loon
24 Goldfinch	31 Brown Creeper

Total number of migrants seen, 55.

SILENE CONICA IN MICHIGAN.

BY J. A. NIEUWLAND.

As far as I have been able to find only two places in the region covered by Gray's and Britton's manuals have reported the presence of the European plant adventive to America. I have found the plant so well established near Stephenville, Mich., (Berrien Co.), as to be likely to become a more or less noxious weed. It was growing abundantly in rather sandy well cultivated strawberry patches and along fences, about one and one half miles northwest of the village, about one mile from Lake Michigan itself and about five and one half or six miles south of St. Joseph, Michigan. As the plant at first sight seemed new to me I had gathered considerable material at the time. The specimens collected are No. 2713 of the University herbarium. Judging from the close aggregation of the plants, and the apparent health of the specimens, it would seem to be well adapted to the soil, and may become a weed that the farmer will have to reckon with if unchecked.

SYNONYMY ALTERATIONS.

Laciniaria scariosa media to replace *L. scariosa intermedia* (*Liatris intermedia* Lindl. Bot. Reg. t. 948).

Laciniaria scariosa Nieuwlandii septentrionalis to replace *L. scariosa Nieuwlandii borealis* (*Liatris borealis* Nutt. in Paxt. Mag. v.t. 27). And *Laciniaria scariosa borealis* (Nutt.), l. c., to replace *L. scariosa virginiana*.

Leeds, North Dakota.

May number of Midland Naturalist was published April 18, 1912.

Pages 29-35 incl. with Plate I. of Vol. II., Am. Mid. Naturalist were published as a preliminary separate Jan. 31, 1911.